

INSTRUCTOR'S SOLUTIONS MANUAL

INTRODUCTION TO MATHEMATICAL STATISTICS SEVENTH EDITION

Robert Hogg

University of Iowa


Joseph McKean

Western Michigan University

Allen Craig

PEARSON

Boston Columbus Indianapolis New York San Francisco Upper Saddle River
Amsterdam Cape Town Dubai London Madrid Milan Munich Paris Montreal Toronto
Delhi Mexico City Sao Paulo Sydney Hong Kong Seoul Singapore Taipei Tokyo



The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expressed or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson from electronic files supplied by the author.

Copyright © 2013, 2005, 1995 Pearson Education, Inc.
Publishing as Pearson, 75 Arlington Street, Boston, MA 02116.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.

ISBN-13: 978-0-321-79565-6
ISBN-10: 0-321-79565-2

www.pearsonhighered.com

PEARSON

Contents

1	Probability and Distributions	1
2	Multivariate Distributions	11
3	Some Special Distributions	19
4	Some Elementary Statistical Inferences	31
5	Consistency and Limiting Distributions	49
6	Maximum Likelihood Methods	53
7	Sufficiency	65
8	Optimal Tests of Hypotheses	77
9	Inferences about Normal Models	83
10	Nonparametric and Robust Statistics	93
11	Bayesian Statistics	103

Chapter 1

Probability and Distributions

1.2.1 Part (c): $C_1 \cap C_2 = \{(x, y) : 1 < x < 2, 1 < y < 2\}$.

1.2.3 $C_1 \cap C_2 = \{\text{mary, mray}\}$.

1.2.6 $C_k = \{x : 1/k \leq x \leq 1 - (1/k)\}$.

1.2.7 $C_k = \{(x, y) : 0 \leq x \leq 1/k, 0 \leq y \leq 1/k\}$.

1.2.8 $\lim_{k \rightarrow \infty} C_k = \{x : 0 < x < 3\}$. Note: neither the number 0 nor the number 3 is in any of the sets C_k , $k = 1, 2, 3, \dots$

1.2.9 Part (b): $\lim_{k \rightarrow \infty} C_k = \phi$, because no point is in all the sets C_k , $k = 1, 2, 3, \dots$

1.2.11 Because $f(x) = 0$ when $1 \leq x < 10$,

$$Q(C_3) = \int_0^{10} f(x) dx = \int_0^1 6x(1-x) dx = 1.$$

1.2.13 Part (c): Draw the region C carefully, noting that $x < 2/3$ because $3x/2 < 1$.
Thus

$$Q(C) = \int_0^{2/3} \left[\int_{x/2}^{3x/2} dy \right] dx = \int_0^{2/3} x dx = 2/9.$$

1.2.16 Note that

$$25 = Q(\mathcal{C}) = Q(C_1) + Q(C_2) - Q(C_1 \cap C_2) = 19 + 16 - Q(C_1 \cap C_2).$$

Hence, $Q(C_1 \cap C_2) = 10$.

1.2.17 By studying a Venn diagram with 3 intersecting sets, it should be true that

$$11 \geq 8 + 6 + 5 - 3 - 2 - 1 = 13.$$

It is not, and the accuracy of the report should be questioned.

1.3.3

$$P(\mathcal{C}) = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots = \frac{1/2}{1 - (1/2)} = 1.$$

1.3.6

$$P(\mathcal{C}) = \int_{-\infty}^{\infty} e^{-|x|} dx = \int_{-\infty}^0 e^x dx + \int_0^{\infty} e^{-x} dx = 2 \neq 1.$$

We must multiply by 1/2.

1.3.8

$$P(C_1^c \cup C_2^c) = P[(C_1 \cap C_2)^c] = P(\mathcal{C}) = 1,$$

because $C_1 \cap C_2 = \phi$ and $\phi^c = \mathcal{C}$.

1.3.11 The probability that he does not win a prize is

$$\binom{990}{5} / \binom{1000}{5}.$$

1.3.13 Part (a): We must have 3 even or one even, 2 odd to have an even sum. Hence, the answer is

$$\frac{\binom{10}{3}\binom{10}{0}}{\binom{20}{3}} + \frac{\binom{10}{1}\binom{10}{2}}{\binom{20}{3}}.$$

1.3.14 There are 5 mutual exclusive ways this can happen: two “ones”, two “twos”, two “threes”, two “reds”, two “blues.” The sum of the corresponding probabilities is

$$\frac{\binom{2}{2}\binom{6}{0} + \binom{2}{2}\binom{6}{0} + \binom{2}{2}\binom{6}{0} + \binom{5}{2}\binom{3}{0} + \binom{3}{2}\binom{5}{0}}{\binom{8}{2}}.$$

1.3.15

$$\begin{aligned} \text{(a)} \quad & 1 - \frac{\binom{48}{5}\binom{2}{0}}{\binom{50}{5}} \\ \text{(b)} \quad & 1 - \frac{\binom{48}{n}\binom{2}{0}}{\binom{50}{n}} \geq \frac{1}{2}, \text{ Solve for } n. \end{aligned}$$

1.3.20 Choose an integer $n_0 > \max\{a^{-1}, (1-a)^{-1}\}$. Then $\{a\} = \bigcap_{n=n_0}^{\infty} (a - \frac{1}{n}, a + \frac{1}{n})$. Hence by (1.3.10),

$$P(\{a\}) = \lim_{n \rightarrow \infty} P\left[\left(a - \frac{1}{n}, a + \frac{1}{n}\right)\right] = \frac{2}{n} = 0.$$

1.4.2

$$P[(C_1 \cap C_2 \cap C_3) \cap C_4] = P[C_4|C_1 \cap C_2 \cap C_3]P(C_1 \cap C_2 \cap C_3),$$

and so forth. That is, write the last factor as

$$P[(C_1 \cap C_2) \cap C_3] = P[C_3|C_1 \cap C_2]P(C_1 \cap C_2).$$

1.4.5

$$\frac{\left[\binom{4}{3} \binom{48}{10} + \binom{4}{4} \binom{48}{9} \right] / \binom{52}{13}}{\left[\binom{4}{2} \binom{48}{11} + \binom{4}{3} \binom{48}{10} + \binom{4}{4} \binom{48}{9} \right] / \binom{52}{13}}.$$

1.4.10

$$P(C_1|C) = \frac{(2/3)(3/10)}{(2/3)(3/10) + (1/3)(8/10)} = \frac{3}{7} < \frac{2}{3} = P(C_1).$$

1.4.12 Part (c):

$$\begin{aligned} P(C_1 \cup C_2^c) &= 1 - P[(C_1 \cup C_2^c)^c] = 1 - P(C_1^* \cap C_2) \\ &= 1 - (0.4)(0.3) = 0.88. \end{aligned}$$

1.4.14 Part (d):

$$1 - (0.3)^2(0.1)(0.6).$$

1.4.16 $1 - P(TT) = 1 - (1/2)(1/2) = 3/4$, assuming independence and that H and T are equilikely.

1.4.19 Let C be the complement of the event; i.e., C equals at most 3 draws to get the first spade.

$$\begin{aligned} \text{(a)} \quad P(C) &= \frac{1}{4} + \frac{3}{4} \frac{1}{4} + \left(\frac{3}{4}\right)^2 \frac{1}{4}. \\ \text{(b)} \quad P(C) &= \frac{1}{4} + \frac{13}{51} \frac{39}{52} + \frac{13}{50} \frac{38}{51} \frac{39}{52}. \end{aligned}$$

1.4.22 The probability that A wins is $\sum_{n=0}^{\infty} \left(\frac{5}{6} \frac{4}{6}\right)^n \frac{1}{6} = \frac{3}{8}$.

1.4.27 Let Y denote the bulb is yellow and let T_1 and T_2 denote bags of the first and second types, respectively.

(a)

$$P(Y) = P(Y|T_1)P(T_1) + P(Y|T_2)P(T_2) = \frac{20}{25} \cdot 6 + \frac{10}{25} \cdot 4.$$

(b)

$$P(T_1|Y) = \frac{P(Y|T_1)P(T_1)}{P(Y)}.$$

1.4.30 Suppose without loss of generality that the prize is behind curtain 1. Condition on the event that the contestant switches. If the contestant chooses curtain 2 then she wins, (In this case Monte cannot choose curtain 1, so he must choose curtain 3 and, hence, the contestant switches to curtain 1). Likewise, in the case the contestant chooses curtain 3. If the contestant chooses curtain 1, she loses. Therefore the conditional probability that she wins is $\frac{2}{3}$.

1.4.31 (1) The probability is $1 - \left(\frac{5}{6}\right)^4$.

$$\text{(2) The probability is } 1 - \left[\left(\frac{5}{6}\right)^2 + \frac{10}{36} \right]^{24}.$$

1.5.2 Part (a):

$$c[(2/3) + (2/3)^2 + (2/3)^3 + \dots] = \frac{c(2/3)}{1 - (2/3)} = 2c = 1,$$

so $c = 1/2$.

1.5.5 Part (a):

$$p(x) = \begin{cases} \frac{\binom{13}{x}\binom{39}{5-x}}{\binom{52}{5}} & x = 0, 1, \dots, 5 \\ 0 & \text{elsewhere.} \end{cases}$$

1.5.9 Part (b):

$$\sum_{x=1}^{50} x/5050 = \frac{50(51)}{2(5050)} = \frac{51}{202}.$$

1.5.10 For Part (c): Let $C_n = \{X \leq n\}$. Then $C_n \subset C_{n+1}$ and $\cup_n C_n = R$. Hence, $\lim_{n \rightarrow \infty} F(n) = 1$. Let $\epsilon > 0$ be given. Choose n_0 such that $n \geq n_0$ implies $1 - F(n) < \epsilon$. Then if $x \geq n_0$, $1 - F(x) \leq 1 - F(n_0) < \epsilon$.

1.6.2 Part (a):

$$p(x) = \frac{\binom{9}{x-1}}{\binom{10}{x-1}} \frac{1}{11-x} = \frac{1}{10}, \quad x = 1, 2, \dots, 10.$$

1.6.3

$$(a) \quad p(x) = \left(\frac{5}{6}\right)^{x-1} \left(\frac{1}{6}\right), \quad x = 1, 2, 3, \dots$$

$$(b) \quad \sum_{x=1}^{\infty} \left(\frac{5}{6}\right)^{x-1} \left(\frac{1}{6}\right) = \frac{1/6}{1 - (25/36)} = \frac{6}{11}.$$

1.6.8 $\mathcal{D}_y = \{1, 2^3, 3^3, \dots\}$. The pmf of Y is

$$p(y) = \left(\frac{1}{2}\right)^{y^{1/3}}, \quad y \in \mathcal{D}_y.$$

1.7.1 If $\sqrt{x} < 10$ then

$$F(x) = P[X(c) = c^2 \leq x] = P(c \leq \sqrt{x}) = \int_0^{\sqrt{x}} \frac{1}{10} dz = \frac{\sqrt{x}}{10}.$$

Thus

$$f(x) = F'(x) = \begin{cases} \frac{1}{20\sqrt{x}} & 0 < x < 100 \\ 0 & \text{elsewhere.} \end{cases}$$

1.7.2

$$C_2 \subset C_1^c \Rightarrow P(C_2) \leq P(C_1^c) = 1 - (3/8) = 5/8.$$

1.7.4 Among other characteristics,

$$\int_{-\infty}^{\infty} \frac{1}{\pi(1+x^2)} dx = \frac{1}{\pi} \arctan x \Big|_{-\infty}^{\infty} = \frac{1}{\pi} \left[\frac{\pi}{2} - \left(-\frac{\pi}{2} \right) \right] = 1.$$

1.7.6 Part (b):

$$\begin{aligned} P(X^2 < 9) &= P(-3 < X < 3) = \int_{-2}^3 \frac{x+2}{19} dx \\ &= \frac{1}{18} \left[\frac{x^2}{2} + 2x \right]_{-2}^3 = \frac{1}{18} \left[\frac{21}{2} - (-2) \right] = \frac{25}{36}. \end{aligned}$$

1.7.8 Part (c):

$$f'(x) = \frac{1}{2} 2x e^{-x} = 0;$$

hence, $x = 2$ is the mode because it maximizes $f(x)$.

1.7.9 Part (b):

$$\int_0^m 3x^2 dx = \frac{1}{2};$$

hence, $m^3 = 2^{-1}$ and $m = (1/2)^{1/3}$.

1.7.10

$$\int_0^{\xi_{0.2}} 4x^3 dx = 0.2;$$

hence, $\xi_{0.2}^4 = 0.2$ and $\xi_{0.2} = 0.2^{1/4}$.

1.7.13 $x = 1$ is the mode because for $0 < x < \infty$ because

$$\begin{aligned} f(x) &= F'(x) = e^{-x} - e^{-x} + x e^{-x} = x e^{-x} \\ f'(x) &= -x e^{-x} + e^{-x} = 0, \end{aligned}$$

and $f'(1) = 0$.

1.7.16 Since $\Delta > 0$

$$X > z \Rightarrow Y = X + \Delta > z.$$

Hence, $P(X > z) \leq P(Y > z)$.

1.7.19 Since $f(x)$ is symmetric about 0, $\xi_{.25} < 0$. So we need to solve,

$$\int_{-2}^{\xi_{.25}} \left(-\frac{x}{4} \right) dx = .25.$$

The solution is $\xi_{.25} = -\sqrt{2}$.

1.7.20 For $0 < y < 27$,

$$x = y^{1/3}, \quad \frac{dx}{dy} = \frac{1}{3}y^{-2/3}$$

$$g(y) = \frac{1}{3y^{2/3}} \frac{y^{2/3}}{9} = \frac{1}{27}.$$

1.7.22

$$f(x) = \frac{1}{\pi}, \quad -\frac{\pi}{2} < x < \frac{\pi}{2}.$$

$$x = \arctan y, \quad \frac{dx}{dy} = \frac{1}{1+y^2}, \quad -\infty < y < \infty.$$

$$g(y) = \frac{1}{\pi} \frac{1}{1+y^2}, \quad -\infty < y < \infty.$$

1.7.23

$$G(y) = P(-2 \log X^4 \leq y) = P(X \geq e^{-y/8}) = \int_{e^{-y/8}}^1 4x^3 dx = 1 - e^{-y/2}, \quad 0 < y < \infty$$

$$g(y) = G'(y) = \begin{cases} e^{-y/2} & 0 < y < \infty \\ 0 & \text{elsewhere.} \end{cases}$$

1.7.24

$$G(y) = P(X^2 \leq y) = P(-\sqrt{y} \leq X \leq \sqrt{y})$$

$$= \begin{cases} \int_{-\sqrt{y}}^{\sqrt{y}} \frac{1}{3} dx = \frac{2\sqrt{y}}{3} & 0 \leq y < 1 \\ \int_{-1}^{\sqrt{y}} \frac{1}{3} dx = \frac{\sqrt{y}}{3} + \frac{1}{3} & 1 \leq y < 4 \end{cases}$$

$$g(y) = \begin{cases} \frac{1}{3\sqrt{y}} & 0 \leq y < 1 \\ \frac{1}{6\sqrt{y}} & 1 \leq y < 4 \\ 0 & \text{elsewhere.} \end{cases}$$

1.8.4

$$E(1/X) = \sum_{x=51}^{100} \frac{1}{x} \frac{1}{50}.$$

The latter sum is bounded by the two integrals

$$\int_{51}^{101} \frac{1}{x} dx \quad \text{and} \quad \int_{50}^{100} \frac{1}{x} dx.$$

An appropriate approximation might be

$$\frac{1}{50} \int_{50.5}^{101.5} \frac{1}{x} dx = \frac{1}{50} (\log 101.5 - \log 50.5).$$

1.8.6

$$E[X(1-X)] = \int_0^1 x(1-x)3x^2 dx.$$

1.8.8 When $1 < y < \infty$

$$G(y) = P(1/X \leq y) = P(X \geq 1/y) = \int_{1/y}^1 2x dx = 1 - \frac{1}{y^2}$$

$$g(y) = \frac{2}{y^3}$$

$$E(Y) = \int_1^\infty y \frac{2}{y^3} dy = 2, \text{ which equals } \int_0^1 (1/x)2x dx.$$

1.8.10 The expectation of X does not exist because

$$E(|X|) = \frac{2}{\pi} \int_0^\infty \frac{x}{1+x^2} dx = \frac{1}{\pi} \int_1^\infty \frac{1}{u} du = \infty,$$

where the change of variable $u = 1 + x^2$ was used.

1.9.2

$$M(t) = \sum_{x=1}^\infty \left(\frac{e^t}{2}\right)^x = \frac{e^t/2}{1-(e^t/2)}, \quad e^t/2 < 1.$$

Find $E(X) = M'(0)$ and $\text{Var}(X) = M''(0) - [M'(0)]^2$.

1.9.4

$$0 \leq \text{var}(X) = E(X^2) - [E(X)]^2.$$

1.9.6

$$E \left[\left(\frac{X - \mu}{\sigma} \right)^2 \right] = \frac{1}{\sigma^2} \sigma^2 = 1.$$

1.9.8

$$K(b) = E[(X-b)^2] = E(X^2) - 2bE(X) + b^2$$

$$K'(b) = -2E(X) + 2b = 0 \Rightarrow b = E(X).$$

1.9.11 For a continuous type random variable,

$$K(t) = \int_{-\infty}^\infty t^x f(x) dx.$$

$$K'(t) = \int_{-\infty}^\infty x t^{x-1} f(x) dx \Rightarrow K'(1) = E(X).$$

$$K''(t) = \int_{-\infty}^\infty x(x-1)t^{x-2} f(x) dx \Rightarrow K''(1) = E[X(X-1)];$$

and so forth.

1.9.12

$$\begin{aligned}
 3 &= E(X - 7) \Rightarrow E(X) = 10 = \mu. \\
 11 &= E[(X - 7)^2] = E(X^2) - 14E(X) + 49 = E(X^2) - 91 \\
 &\Rightarrow E(X^2) = 102 \text{ and } \text{var}(X) = 102 - 100 = 2. \\
 15 &= E[(X - 7)^3]. \text{ Expand } (X - 7)^3 \text{ and continue.}
 \end{aligned}$$

1.9.16

$$\begin{aligned}
 E(X) &= 0 \Rightarrow \text{var}(X) = E(X^2) = 2p. \\
 E(X^4) &= 2p \Rightarrow \text{kurtosis} = 2p/4p^2 = 1/2p.
 \end{aligned}$$

1.9.17

$$\begin{aligned}
 \psi'(t) &= M'(t)/M(t) \Rightarrow \psi'(0) = M'(0)/M(0) = E(X). \\
 \psi''(t) &= \frac{M(t)M''(t) - M'(t)M'(t)}{[M(t)]^2} \\
 \Rightarrow \psi''(0) &= \frac{M(0)M''(0) - M'(0)M'(0)}{[M(0)]^2} = M''(0) - [M'(0)]^2 = \text{var}(X).
 \end{aligned}$$

1.9.19

$$M(t) = (1 - t)^{-3} = 1 + 3t + 3 \cdot 4 \frac{t^2}{2!} + 3 \cdot 4 \cdot 5 \frac{t^3}{3!} + \dots$$

Considering the coefficient of $t^r/r!$, we have

$$E(X^r) = 3 \cdot 4 \cdot 5 \cdots (r + 2), \quad r = 1, 2, 3, \dots$$

1.9.20 Integrating the parts with $u = 1 - F(x)$, $dv = dx$, we get

$$\{[1 - F(x)]x\}_0^b - \int_0^b x[-f(x)] dx = \int_0^b xf(x) dx = E(X).$$

1.9.23

$$\begin{aligned}
 E(X) &= \int_0^1 x \frac{1}{4} dx + 0 \cdot \frac{1}{4} + 1 \cdot \frac{1}{2} = \frac{5}{8}. \\
 E(X^2) &= \int_0^1 x^2 \frac{1}{4} dx + 0 \cdot \frac{1}{4} + 1 \cdot \frac{1}{2} = \frac{7}{12}. \\
 \text{var}(X) &= \frac{7}{12} - \left(\frac{5}{8}\right)^2 = \frac{37}{192}.
 \end{aligned}$$

1.9.24

$$E(X) = \int_{-\infty}^{\infty} x[c_1 f_1(x) + \cdots + c_k f_k(x)] dx = \sum_{i=1}^k c_i \mu_i = \mu.$$

Because $\int_{-\infty}^{\infty} (x - \mu)^2 f_i(x) dx = \sigma_i^2 + (\mu_i - \mu)^2$, we have

$$E[(X - \mu)^2] = \sum_{i=1}^k c_i [\sigma_i^2 + (\mu_i - \mu)^2].$$

1.10.2

$$\mu = \int_0^{\infty} x f(x) dx \geq \int_{2\mu}^{\infty} 2\mu f(x) dx = 2\mu P(X > 2\mu).$$

Thus $\frac{1}{2} \geq P(X > 2\mu)$.

1.10.4 If, in Theorem 1.10.2, we take $u(X) = \exp\{tX\}$ and $c = \exp\{ta\}$, we have

$$P(\exp\{tX\} \geq \exp\{ta\}) \leq M(t) \exp\{-ta\}.$$

If $t > 0$, the events $\exp\{tX\} \geq \exp\{ta\}$ and $X \geq a$ are equivalent. If $t < 0$, the events $\exp\{tX\} \geq \exp\{ta\}$ and $X \leq a$ are equivalent.

1.10.5 We have $P(X \geq 1) \leq [1 - \exp\{-2t\}]/2t$ for all $0 < t < \infty$, and $P(X \leq -1) \leq [\exp\{2t\} - 1]/2t$ for all $-\infty < t < 0$. Each of these bounds has the limit 0 as $t \rightarrow \infty$ and $t \rightarrow -\infty$, respectively.